

OP data on CD for mean opacities and radiative accelerations

M.J. Seaton

Department of Physics and Astronomy, University College London, London WC1E 6BT

Accepted XXX. Received XXX; in original form XXX

ABSTRACT

All monochromatic opacity data from the Opacity Project (OP), together with all codes required for the calculation of mean opacities and radiative accelerations for any required chemical mixture, temperature and mass-density, are being put on a 700 Mb CD which will be made generally available. The present paper gives a concise summary of the contents of the CD. More complete documentation will be provided on the CD itself.

Key words: atomic processes – radiative transfer – stars: interiors.

1 INTRODUCTION

Methods used by the Opacity Project (OP) for the calculation of mean opacities (Planck and Rosseland) were described in Paper I (Seaton *et al.*, 1994); further work on the inclusion of contributions from inner-shell transitions were discussed in Paper II (Badnell and Seaton, 2003) for the six-element mix of Iglesias and Rogers (1995); comparisons with OPAL (see Rogers & Iglesias, 1992) for that mix in Paper III (Seaton and Badnell, 2004); and up-dated results for all elements, with inclusion of inner-shell contributions, in Paper IV (Badnell *et al.*, 2004). The OP approach to the calculation of radiative accelerations is discussed by Seaton (1997).

The basic monochromatic opacity data for the 17 elements H, He, C, N, O, Ne, Na, Mg, Al, Si, S, Ar, Ca, Cr, Mn, Fe and Ni occupy 665 Mb as **gzip** formatted files, or 868 Mb as binary files. Some improvements have been made in the frequency-mesh which is used (see section 2.2.1 below) which lead to improved accuracies in the numerical integrations required to obtain mean opacities and radiative accelerations. It is found that the integrations for Rosseland-means never give numerical errors larger than 0.1%. Improvements have been made in the codes used to obtain mean opacities and accelerations for any required chemical mixture, interpolated to any required values of temperature and mass-density. The interpolated Rosseland mean opacities are shown to be numerically correct to better than 1% for all practical purposes.

It is our intention to make all OP data generally available, both on a 700 Mb CD which will be described in the present paper and on a web OPserver which will be described by Mendoza *et al.* (2004).

The CD contains all data required for the calculation of OP mean opacities and radiative accelerations, together

with full documentation and all required codes, and is available on request¹. It contains a **tar** file of compressed (**gzip**) files. The file **OPCD.ps.gz** provides full documentation, including information about processing of all other data on the **tar** file.

The user is urged to read the file **OPCD.ps** *before* decompressing any other files.

The present paper gives a concise summary of the contents of the CD. Full documentation is given in the file **OPCD.ps**. The CD includes input and output files for some test runs.

2 BASIC DEFINITIONS

2.1 Monochromatic opacities

The monochromatic opacity cross-section for element k is $\sigma_k(u)$ where $u = h\nu/(k_B T)$ and k_B is Boltzmann's constant. The correction-factor for stimulated emission, $[1 - \exp(-u)]$, is *not* included² in the definition of σ_k . For a mixture of elements let element k have abundance by number-fraction of f_k , $\sum_k f_k = 1$. The cross-section for the mixture is

$$\sigma(u) = \sum_k f_k \sigma_k(u). \quad (1)$$

2.2 The Rosseland mean

The Rosseland-mean cross-section is

¹ Requests to: Claude Zeippen (Claude.Zeippen@obspm.fr); or Anil Pradhan (pradhan.1@osu.edu)

² We here consider the quantity actually tabulated in the OP archives. A number of previous papers have defined σ_k *with* inclusion of the stimulated emission factor.

$$\frac{1}{\sigma_R} = \int_0^\infty \frac{F(u)}{\sigma(u)[1 - \exp(-u)]} du \quad (2)$$

where

$$F(u) = [15/(4\pi^4)]u^4 \exp(-u)/[1 - \exp(-u)]^2. \quad (3)$$

The Rosseland-mean opacity, per unit mass, is $\kappa_R = \sigma_R/\mu$ where μ is mean atomic weight.

Numerical values for cross-sections σ will be given in atomic units (a_0^2) and values for opacities κ_R in cgs units ($\text{cm}^2 \text{g}^{-1}$).

2.2.1 The frequency variable

It is convenient to use the frequency-variable³

$$v(u) = \int_0^u \frac{F(u)}{[1 - \exp(-u)]} du \quad (4)$$

giving

$$\frac{1}{\sigma_R} = \int_0^{v_\infty} \frac{1}{\sigma(u)} dv \quad (5)$$

where $v_\infty = v(u \rightarrow \infty)$

2.3 Radiative accelerations

The radiative acceleration for element k in a stellar interior is

$$g_{\text{rad}} = (1/c)(\mu/\mu_k)\mathcal{F}\kappa_R\gamma_k \quad (6)$$

where c is the speed of light, μ_k the atomic mass of element k and

$$\mathcal{F} = \pi B(T_{\text{eff}})(R_\star/r)^2 \quad (7)$$

where T_{eff} is the effective temperature,

$$B(T) = 2(\pi k_B T)^4 / (15c^2 h^3), \quad (8)$$

$R_\star$ is the radius of the star and r the distance from the centre of the star. In (6),

$$\gamma_k = \int \frac{\sigma_k^{\text{mta}}}{\sigma} dv \quad (9)$$

where σ_k^{mta} is a cross-section for momentum-transfer to the atom,

$$\sigma_k^{\text{mta}} = \sigma_k(u)[1 - \exp(-u)] - a_k(u). \quad (10)$$

In (10), $a_k(u)$ is a correction factor to take out contributions from scattering processes and momentum transfer to electrons in photo-ionization processes.

Values of g_{rad} are given in cgs units (cm s^{-2}). The quantity γ_k is dimensionless.

One is usually interested in the case in which the abundance of the selected element k varies with depth in a star due to the operation of diffusion processes. The codes allow for the possibility of varying the abundance of k by a factor χ , keeping the relative abundances of all other elements unchanged.

³ Some previous papers have taken σ to be defined *with* inclusion of the stimulated emission factor, and v to be defined by $v = \int F(u) du$.

3 ARCHIVED DATA FILES

All data are tabulated on a mesh of values of temperature T and electron-density N_e . The indices **ite** and **jne** are defined by

$$\text{ite} = 40 \times \log(T), \quad \text{jne} = 4 \times \log(N_e). \quad (11)$$

All data on the CD are with the “m” mesh, even values of **ite** and **jne**, $\Delta \text{ite} = \Delta \text{jne} = 2$.

The cross-sections are tabulated for **ntot** = 10000 equally-spaced values of v . The σ_k data are in files **mzz.ttt** and the a_k are in files **azz.ttt** where **zz** is a two-digit number specifying nuclear charge (**zz** = 01 for H) and **ttt** is a three-digit number specifying **ite**. Data for all **jne** included are contained in each files **mzz.ttt** or **azz.ttt**. Files **mzz.mesh** give values of u for the tabulated values of v .

The OP codes use unformatted (binary) data files, for economy in storage and speed of input/output. However, the CD contains files which are formatted and compressed (**gzip**) for ease of transfer between machines. The procedures to be used for converting to binary are described in the file **OPCD.ps**.

Some summary data, described in the documentation, are given in files **mzz.index**, **mzz.ion**, **mzz.smry**.

4 CODES FOR ROSSELAND-MEANS

4.1 The code **mixv.f**

The code **mixv.f** reads a list of abundances f_k for a mixture, obtains the cross-sections σ and produces an output file with values of mass-density ρ and Rosseland means κ_R for all (T, N_e) mesh points.

4.2 The code **opfit.f**

The code **opfit.f** reads an output file from **mixv.f** and a list of values of ($\log(T), \log(\rho)$) for a stellar model and produces a file giving values of

$$\log(\kappa_R), \quad \partial \log(\kappa_R) / \partial \log(T), \quad \partial \log(\kappa_R) / \partial \log(\rho). \quad (12)$$

The code uses refined techniques of bi-cubic spline interpolations. It has an optional facility (rarely required) to smooth the data before doing the interpolations, and an optional feature to produce output tables in OPAL format.

4.3 The code **mx.f**

The code **mx.f** produces output similar to that from **opfit.f** in a one-step process, using less refined bicubic interpolation techniques. It has a facility that abundances can be changed at any point during a run for a particular stellar model.

4.4 The code **mixz.f**

The code **mixz.f** produces monochromatic opacities for any specified mixture of metals. The output files are stored as a “fictitious” metal, **mzz.ttt** with **zz** a two-digit number greater than 28 (data for Nickel are stored in files **m28.ttt**).

4.5 The code `mxz.f`

The code `mxz.f` is similar to `mx.f` but uses the files for a metal-mixture from `mixz.f`. It has a facility that, at each depth point, the values of X and Z can be changed.

4.6 Choice of mixture codes

Use of `mixv.f` plus `opfit.f` can be convenient if the same mixture is to be used for many stellar models. The use of splines in `opfit.f` ensures that the derivatives in (12) are continuous. It is shown in Paper IV that, for all cases likely to be of practical interest, the interpolations using `opfit.f` do not introduce any errors as large as 1%.

Use of `mx.f` is particularly convenient if there are changes in composition depending on depth in a stellar model. In occasional difficult cases use of `mx.f` can give a lack of smoothness in the derivatives.

The test runs of `mixv.f`, `opfit.f` and `mx.f` are for S92 abundances (see Paper I), $\log(T) = 6.0$ to 7.3 and $\log(R) = -1.75$, roughly corresponding to conditions in the solar radiative interior. Use of `mx.f` gives values of $\log(\kappa_R)$ and $\partial \log(\kappa_R)/\partial \log(T)$ which appear to be smooth and in agreement with values from `opfit.f` but there is some apparent lack of smoothness in $\partial \log(\kappa_R)/\partial \log(\rho)$ from `mx.f`, as shown on Figure 1.

The codes `mixz.f` and `mxz.f` were used in Paper IV for the case a more realistic solar model in which, below the base of the convection zone, X and Z varied continuously as functions of depth.

5 CODES FOR RADIATIVE ACCELERATIONS

5.1 The code `accv.f`

The code `accv.f` has a structure similar to that of `mixv.f`. It reads a list of abundances and specification of the selected element k , reads data specifying a range of values of the abundance-multiplier χ , reads the files `mzz.ttt` and `azz.ttt` and produces an output file giving, for each (T, N_e) mesh point, values of

$$\kappa_R, \partial \kappa_R / \partial \chi, \gamma, \partial \gamma / \partial \chi \quad (13)$$

for each value of χ .

5.2 The code `accfit.f`

The code `accfit.f` is similar to `opfit.f` using bi-cubic splines. It reads a file output from `accv.f`, a list of values of χ , the effective temperature T_{eff} for a stellar model and a list of values of $\log(T)$, $\log(\rho)$ and r/R_* . It produces a file giving value of

$$\log(\kappa_R), \log(\gamma), \log(g_{\text{rad}}) \quad (14)$$

as functions of χ .

5.3 The code `ax.f`

The code `ax.f` has a structure similar to that of `mx.f`. Its use is convenient when the mixture of elements depends on depth in a stellar model.

5.4 Choice of codes for accelerations

Use of `accfit.f` provides the most accurate interpolations. The code `ax.f` can be useful for cases in which abundances vary with depth.

The test case is for Argon in a stellar model with $T_{\text{eff}} = 10^4 \text{K}$. Figure 2 shows g_{rad} for $\chi = 0.01, 1.00$ and 100 . The deep minimum in the vicinity of $\log(T) = 5.45$ occurs when the Argon is going through a Neon-like ionization stage. It is seen that g_{rad} from `ax.f` shows some lack of smoothness for the rather extreme case of $\chi = 0.01$

REFERENCES

- Badnell N.R., Seaton M.J., 2003. J. Phys. B, 36,4367 (Paper II)
 Badnell N.R., Bautista M.A., Butler K., Delahaye F., Mendoza C., Palmeri P., Zeippen C.J., Seaton M.J., 2004. MNRAS, to be submitted (Paper IV)
 Iglesias C.A., Rogers, F.J., 1995, ApJ 443, 469
 Mendoza, C. *et al.*, 2004, MNRAS, to be submitted
 Rogers, F.J., Iglesias, C.A., 1992. ApJS 79,507, 1992
 Seaton, M.J., 1997, MNRAS, 289, 700
 Seaton, M.J., Badnell, N.R., 2004, MNRAS, in press (Paper III)
 Seaton, M.J., Yu Yan, Mihalas D., Pradhan, A.K., 1994, MNRAS, 266, 805 (Paper I)




Figure 1 $\partial \log(\kappa_R)/\partial \log(\rho)$ for a model with S92 abundances and $\log(R) = -1.75$. Results obtained using the codes `opfit.f` and `mx.f`.




Figure 2 Radiative accelerations for Ar in a model with $T_{\text{eff}} = 10^4 \text{K}$. Results from codes `accv.f` and `ax.f` with abundance multipliers $\chi = 0.01, 1.0$ and 100 .